

## References and further reading

- Ceyhan D, Gucyetmez Topal B.Ceyhan D, et al. [An 18-Month-Old Child with Infantile Pompe Disease: Oral Signs.](#) Case Rep Dent. 2017;2017:5685941. doi: 10.1155/2017/5685941.
- De Gijt JP, Van Capelle CI, Oosterhuis JW, Van Der Ploeg AT, Van Der Wal KG.De Gijt JP, et al. [Gingival overgrowth in Pompe disease: a case report.](#) J Oral Maxillofac Surg. 2011 Aug;69(8):2186-90. doi: 10.1016/j.joms.2011.03.070.
- Drążewski D, Grzymisławska M, Korybalska K, Czepulis N, Grzymisławski M, Witowski J, Surdacka A.Drażewski D, et al. [Oral Health Status of Patients with Lysosomal Storage Diseases in Poland.](#) Int J Environ Res Public Health. 2017 Mar 9;14(3):281. doi: 10.3390/ijerph14030281.
- Galeotti A, De Rosa S, Uomo R, Dionisi-Vici C, Deodato F, Taurisano R, Olivieri G, Festa P.Galeotti A, et al. [Orofacial features and pediatric dentistry in the long-term management of Infantile Pompe Disease children.](#) Orphanet J Rare Dis. 2020 Nov 23;15(1):329. doi: 10.1186/s13023-020-01615-1.
- Van Gelder CM, van Capelle CI, Ebbink BJ, Moor-van Nugteren I, Van Den Hout JM, Hakkesteegt MM, Van Doorn PA, De Coo IF, Reuser AJ, De Gier HH, Van Der Ploeg AT. [Facial-muscle weakness, speech disorders and dysphagia are common in patients with classic infantile Pompe disease treated with enzyme therapy.](#) J Inherit Metab Dis. 2012 May;35(3):505-11. doi: 10.1007/s10545-011-9404-7.